



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Reports of States and yearly and monthly reports of cities.*

COLORADO—*Denver*.—Month of September, 1896. Estimated population, 150,000. Total deaths, 123, including diphtheria, 1; enteric fever, 28, and phthisis pulmonalis, 22.

ILLINOIS—*Chicago*.—Month of September, 1896. Estimated population, 1,750,000. Total deaths, 1,658, including diphtheria, 93; measles, 2; scarlet fever, 5; enteric fever, 87; whooping cough, 28, and phthisis pulmonalis, 174.

IOWA—*Dubuque*.—Month of September, 1896. Estimated population, 40,000. Total deaths, 22, including diphtheria, 5; enteric fever, 2, and phthisis pulmonalis, 3.

MICHIGAN.—Reports to the State board of health, Lansing, Mich., from 54 observers indicate that during the week ended October 17, 1896, erysipelas and inflammation of the kidney increased, and typhoid fever decreased in area of prevalence. During the week phthisis pulmonalis was reported at 229 places, enteric fever at 67, diphtheria at 34, scarlet fever at 25, whooping cough at 11, and measles at 10 places.

MISSOURI—*Kansas City*.—Month of September, 1896. Estimated population, 165,000. Total deaths, 126, including diphtheria and croup, 7; enteric fever, 3, and phthisis pulmonalis, 5.

*St. Louis*.—Month of September, 1896. Estimated population, 570,000. Total deaths, 682, including diphtheria and croup, 38; whooping cough, 6; enteric fever, 13, and phthisis pulmonalis, 89.

NEBRASKA—*Omaha*.—Month of August, 1896. Population, United States census 1890, 140,452. Total deaths, 77, including diphtheria and croup, 5; enteric fever, 3, and phthisis pulmonalis, 4.

Month of September, 1896. Total deaths, 63, including diphtheria and croup, 6; enteric fever, 3, and phthisis pulmonalis, 5.

NEW JERSEY—*Paterson*.—Month of September, 1896.—Estimated population, 103,828. Total deaths, 137, including scarlet fever, 3; diphtheria and croup, 18; enteric fever, 5, and phthisis pulmonalis, 5.

UTAH—*Salt Lake City*.—Month of September, 1896. Estimated population, 70,000. Total deaths, 50, including enteric fever, 4, and phthisis pulmonalis, 3.